arXiv:1208.2926v1 [math.NT] 14 Aug 2012

A RELATION BETWEEN MULTIPLICITY ONE AND BÖCHERER'S CONJECTURE

Abstract. We show that a weak form of the generalized Böcherer's conjecture implies multiplicity one for Siegel cusp forms of degree 2.

1. Multiplicity one

Let $S_k(\Gamma)$ denote the space of classical holomorphic cusp forms of weight k for $\Gamma = \mathrm{SL}_2(\mathbb{Z})$. Suppose that f_1 and f_2 are two Hecke eigenforms in $S_k(\Gamma)$ such that the Hecke eigenvalues $\lambda_p(f_1)$ and $\lambda_p(f_2)$ are equal for all primes p. Then, it is well known that there exists a constant c such that $f_1 = cf_2$. Indeed, the various Hecke eigenvalues $\lambda_p(f_i)$ determine all the Fourier coefficients of f_i, hence the form f_i itself, up to a common multiple.

The above fact is a special case of the "multiplicity one theorem" for GL_2 due to Jacquet–Langlands. Multiplicity one was extended to the case of automorphic representations on the group GL_n independently by Shalika, Piatetski-Shapiro, and Gelfand–Kazhdan. However, much less is known for automorphic representations on other reductive groups.

In this note, we consider the space[1] $S_k(\Gamma_2)$ of holomorphic Siegel cusp forms of weight k for $\Gamma_2 = \mathrm{Sp}_4(\mathbb{Z})$. These forms give rise to automorphic representations on the group GSp_4, for which multiplicity one remains unknown. In particular, one has the following conjecture.

Conjecture 1.1 (Multiplicity one for Siegel cusp forms of degree 2 and full level)**.** *Let f_1 and f_2 be two Hecke eigenforms in $S_k(\Gamma_2)$ such that for all primes p, we have an equality of Hecke eigenvalues $\lambda_p(f_1) = \lambda_p(f_2)$ and $\lambda_{p^2}(f_1) = \lambda_{p^2}(f_2)$. Then, there exists a constant c such that $f_1 = cf_2$.*

The above conjecture is deep. It does not appear to directly follow from the transfer to GL_4 that was proved in [7] or even from the more general functoriality and classification results proved recently by Arthur. A key reason why things are so complicated is that Hecke eigenforms in $S_k(\Gamma_2)$ are non-generic, and hence do not have a Whittaker expansion like eigenforms on GL_n do.

Remark 1.2. In Conjecture 1.1, one may replace "for all primes" by "for almost all primes" without increasing the difficulty of the problem; see the argument of [10, Thm. 3.1.5]).

2. Böcherer's conjecture

Another deep and famous conjecture for Siegel cusp forms of degree 2 deals with the relation between the central L-values and the Fourier coefficients. Before stating this conjecture, we briefly recall the Fourier expansion for Siegel cusp forms. For any $f \in S_k(\Gamma_2)$, we can write

---

[1] For definitions and background on Siegel cusp forms, see [4].

$$f(Z)=\sum_S a(f,S)e^{2\pi i \mathrm{Tr}(SZ)},$$

where the Fourier coefficients $a(f,S)$ are indexed by matrices S of the form

$$S=\begin{pmatrix} a & b/2 \\ b/2 & c \end{pmatrix},\qquad a,b,c\in\mathbb{Z},\qquad a>0,\qquad \mathrm{disc}(S):=b^2-4ac<0. \tag{1}$$

Equivalently, the Fourier coefficients $a(f,S)$ are indexed by all positive, integral, binary quadratic forms. In fact, using the defining relation for Siegel cusp forms, we can see that

$$a(f,{}^tASA)=a(f,S) \tag{2}$$

for all $A\in\mathrm{SL}_2(\mathbb{Z})$, thus showing that $a(f,S)$ only depends on the $\mathrm{SL}_2(\mathbb{Z})$-equivalence class of the matrix S, or equivalently, only on the proper equivalence class of the associated binary quadratic form.

Let $d>0$ be an integer such that $-d$ is a fundamental discriminant.[2] Put $K=\mathbb{Q}(\sqrt{-d})$ and let Cl_K denote the ideal class group of K. It is a fact going back to Gauss that the $\mathrm{SL}_2(\mathbb{Z})-$equivalence classes of binary quadratic forms of discriminant $-d$ are in natural bijective correspondence with the elements of Cl_K. In view of the comments above, it follows that for any $f\in S_k(\Gamma_2)$ and any $c\in\mathrm{Cl}_K$ the notation $a(f,c)$ makes sense. We define

$$R(f,K)=\sum_{c\in\mathrm{Cl}_K}a(f,c). \tag{3}$$

Now, suppose that f is an eigenform for the local Hecke algebras at all places. Then f gives rise to an irreducible cuspidal automorphic representation π_f of $\mathrm{GSp}_4(\mathbb{A})$; see [6]. The remarkable conjecture below was first made by Böcherer [1].

Conjecture 2.1 (Böcherer's conjecture)**.** *Let $f\in S_k(\Gamma_2)$ be a non-zero Hecke eigenform and π_f the associated automorphic representation. Then there exists a constant c_f depending only on f such that for any imaginary quadratic field $K=\mathbb{Q}(\sqrt{-d})$ with $-d$ a fundamental discriminant, we have*

$$|R(f,K)|^2=c_f\cdot d^{k-1}\cdot w(K)^2\cdot L(1/2,\pi_f\times\chi_{-d}).$$

Above, χ_{-d} is the quadratic Hecke character associated via class field theory to the field $\mathbb{Q}(\sqrt{-d})$, $w(K)$ denotes the number of distinct roots of unity inside K, and $L(s,\pi_f\times\chi_{-d})$ denotes the associated Langlands L-function.

The above conjecture is deep and so far not proven for any Siegel cusp form that is not a lift of some sort. Theoretical evidence for the truth of a refined version of the above conjecture (see [5, (5.4.5)]) was provided in recent work of the author with Kowalski and Tsimerman.

Böcherer's conjecture has been further generalized by various people. We note in particular the papers by Furusawa–Shalika [3], Furusawa–Martin [2] and Prasad–Takloo-Bighash [8]. In these generalizations, one takes a linear combination of the Fourier coefficients $a(f,c)$ with the values taken by an ideal class character of Cl_K

---

[2] Recall that an integer n is a fundamental discriminant if *either* n is a squarefree integer congruent to 1 modulo 4 *or* $n=4m$ where m is a squarefree integer congruent to 2 or 3 modulo 4.

(more generally, a Hecke character on $\mathbb{A}_K^\times$). Partial progress towards such a conjecture has been made in recent work of Furusawa and Martin. Such a formulation is also closely related to the global Gross-Prasad conjecture for $(SO(5), SO(2))$.

We now describe a "weak version" of a specific refinement of Böcherer conjecture. Let d, K, f, π_f be as before. For any character Λ of the finite group Cl_K, we make the definition

$$R(f, K, \Lambda) = \sum_{c \in \mathrm{Cl}_K} a(f, c)\Lambda^{-1}(c). \tag{4}$$

Also, define the theta-series

$$\theta_\Lambda(z) = \sum_{0 \neq \mathfrak{a} \subset O_K} \Lambda(\mathfrak{a}) e^{2\pi i N(\mathfrak{a}) z}.$$

Thus, θ_Λ is a holomorphic modular form of weight 1 and nebentypus $\left(\frac{-d}{*}\right)$ on $\Gamma_0(d)$; it is a cusp form if and only if $\Lambda^2 \neq 1$. The form θ_Λ generates a dihedral automorphic representation of $\mathrm{GL}_2(\mathbb{A})$ which coincides with the automorphic induction of Λ. We let $L(s, \pi_f \times \theta_\Lambda)$ denote the Langlands L-function attached to the Rankin–Selberg convolution of this dihedral representation with π_f. We now state the

Conjecture 2.2 (Generalized Böcherer's conjecture, weak form)**.** *Let $f \in S_k(\Gamma_2)$ be a non-zero Hecke eigenform and π_f the associated automorphic representation. Let $K = \mathbb{Q}(\sqrt{-d})$ with $-d$ a fundamental discriminant, and Λ be a character of the ideal class group of K. Then $R(f, K, \Lambda) \neq 0$ if and only if $L(1/2, \pi_f \times \theta_\Lambda) \neq 0$.*

Remark 2.3. In the *special case* $\Lambda = 1$, the quantity $R(F, K, \Lambda)$ is simply $R(F, K)$ in our earlier notation and the L-function $L(s, \pi_f \times \theta_\Lambda)$ factors as a product $L(s, \pi_f)L(s, \pi_f \times \chi_{-d})$. Thus, Conjecture 2.2 is compatible with Conjecture 2.1.

Conjecture 2.2 is in the spirit of the global Gross-Prasad conjectures since it asserts that the non-vanishing of a (Bessel) period $R(F, K, \Lambda)$ is equivalent to the non-vanishing of the central value of a related L-function. For a somewhat stronger formulation, see [3, Conj. 1.11].

3. The main result

Theorem 1. *Conjecture 2.2 implies Conjecture 1.1.*

Proof. The proof relies on the following fact which was proved in [9]:

Let $0 \neq f \in S_k(\Gamma_2)$. Then there exists a matrix S of the form given by equation (1) *such that* $\mathrm{disc}(S)$ *is a fundamental discriminant and $a(f, S) \neq 0$.*

Now, assume Conjecture 2.2. Suppose f_1 and f_2 are two Hecke eigenforms in $S_k(\Gamma_2)$ such that for all primes p, we have an equality of Hecke eigenvalues $\lambda_p(f_1) = \lambda_p(f_2)$ and $\lambda_{p^2}(f_1) = \lambda_{p^2}(f_2)$. Let $-d < 0$ be a fundamental discriminant such that there exists a matrix S with $\mathrm{disc}(S) = -d$ and $a(f_2, S) \neq 0$. Such a d exists by the fact quoted above. Put $K = \mathbb{Q}(\sqrt{-d})$ and pick a character Λ of Cl_K such that $R(f_2, K, \Lambda) \neq 0$. Now, put $g_1 = f_1 - \frac{R(f_1,K,\Lambda)}{R(f_2,K,\Lambda)} f_2$.

We claim that $g_1 = 0$. Suppose not. Then g_1 and f_2 are two non-zero Hecke eigenforms in $S_k(\Gamma_2)$ such that for all primes p, we have an equality of Hecke eigenvalues $\lambda_p(g_1) = \lambda_p(f_2)$ and $\lambda_{p^2}(g_1) = \lambda_{p^2}(f_2)$. Since the Hecke operators at p and p^2 generate the full Hecke algebra, it follows that $L(s, \pi_{g_1} \times \theta_\Lambda) = L(s, \pi_{f_2} \times \theta_\Lambda)$.

However, by construction, we have $R(f_2, K, \Lambda) \neq 0$ and $R(g_1, K, \Lambda) = 0$. Since we are assuming Conjecture 2.2 holds true, this means that $L(1/2, \pi_{g_1} \times \theta_\Lambda) = 0$ and $L(1/2, \pi_{g_1} \times \theta_\Lambda) \neq 0$. This is a contradiction. Thus $g_1 = 0$. Hence f_1 and f_2 are multiples of each other.

□

References

[1] Siegfried Böcherer. Bemerkungen über die Dirichletreihen von Koecher und Maass. *Mathematica Gottingensis*, 68:36 pp., 1986.

[2] Masaaki Furusawa and Kimball Martin. On central critical values of the degree four L-functions for GSp(4): The fundamental lemma, II. *American J. Math., to appear*, 2010.

[3] Masaaki Furusawa and Joseph A. Shalika. *On central critical values of the degree four L-functions for* GSp(4), volume 782 of *Memoirs of the American Mathematical Society*. American Mathematical Society, Providence, 2003.

[4] Helmut Kligen. *Introductory lectures on Siegel modular forms*, volume 20 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1990.

[5] Emmanuel Kowalski, Abhishek Saha, and Jacob Tsimerman. Local spectral equidistribution for Siegel modular forms and applications. *Compositio Math.* Forthcoming.

[6] Hiro-aki Narita, Ameya Pitale, and Ralf Schmidt. Irreducibility criteria for local and global representations. *Preprint*, 2011.

[7] Ameya Pitale, Abhishek Saha, and Ralf Schmidt. Transfer of Siegel cusp forms of degree 2. *Preprint.* `arXiv:1106.5611`.

[8] Dipendra Prasad and Ramin Takloo-Bighash. Bessel models for GSp(4). *J. Reine Angew. Math.*, 2011:189–243.

[9] Abhishek Saha. Siegel cusp forms of degree 2 are determined by their fundamental Fourier coefficients. *Math. Ann.*, to appear. `arXiv:1106.5110`.

[10] Ralf Schmidt. Iwahori-spherical representations of GSp(4) and Siegel modular forms of degree 2 with square-free level. *J. Math. Soc. Japan*, 57(1):259–293, 2005.